Turkish Journal of Mathematics

Volume 46 | Number 7 Article 22

1-1-2022

(Co)Limit calculations in the category of 2-crossed R-modules

ELİS SOYLU YILMAZ

Follow this and additional works at: https://journals.tubitak.gov.tr/math

 Part of the Mathematics Commons

Recommended Citation
YILMAZ, ELİS SOYLU (2022) "(Co)Limit calculations in the category of 2-crossed R-modules," *Turkish Journal of Mathematics*: Vol. 46: No. 7, Article 22. https://doi.org/10.55730/1300-0098.3308
Available at: https://journals.tubitak.gov.tr/math/vol46/iss7/22

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Turkish Journal of Mathematics

http://journals.tubitak.gov.tr/math/

Research Article

Turk J Math
(2022) 46: 2902 – 2915
© TÜBİTAK
doi:10.55730/1300-0098.3308

(Co)Limit calculations in the category of 2-crossed R-modules

Elis SOYLU YILMAZ*
Department of Mathematics and Computer Science, Science and Art Faculty, Eskişehir Osmangazi University, Eskişehir, Turkey

Received: 19.12.2021 • **Accepted/Published Online:** 21.07.2022 • **Final Version:** 05.09.2022


Abstract: In this work, we obtain how to construct finite limits and colimits for 2-crossed R-Modules over groups denoted with $\mathbf{X_2Mod/R}$. We give direct construction of the pullback object to show that this category has finite products over the terminal object. We also show finite coproducts and (co)completeness.

Key words: 2-crossed module, coproduct, pullback

1. Introduction

Whitehead introduced the crossed module notion for groups in [20] also known as 2-type groups. 2-crossed modules have been represented by Conduché as characterized of 3-types [11]. Baues [5] defined another version of this concept, as quadratic modules. Brown and Gilbert have given an alternative definition known as braided regular crossed modules for 3-type of groups using the automorphisms structures [6]. Brown, Sivera and Higgins [7] constructed the coproduct for crossed modules of groups. In [4] Arvasi and Ulualan have investigated the relationships between various algebraic types like crossed squares, 2-crossed modules, and simplicial groups.

Crossed squares defined in [16] and in [13] and [14] for commutative algebras. In [12] Conduché obtain a 2-crossed module from the mapping cone of a crossed square. Baues introduced quadratic modules by using simplicial groups which can be regarded as a 2-crossed module endowed with nilpotency conditions. For more details on 2-crossed modules see ([1], [21]). Some of functional relations of 2-crossed modules can be diagramatized as ([3], [2], [17]).



Freeness and coproducts are fundamental tools to construct tensor product for crossed complexes. Freeness of quadratic modules in [2] and 2-crossed modules are given in [18]. In [9] construct the coproduct in

*Correspondence: esoylu@ogu.edu.tr
2010 *AMS Mathematics Subject Classification:* 18A30, 18D35, 20J15

This work is licensed under a Creative Commons Attribution 4.0 International License.

$\mathbf{X_2Mod}/(\mathbf{M} \longrightarrow \mathbf{N})$ that is they fix the precrossed module at the end. Also some colimit theorems and calculations can be seen in [8], [10], [15], and [19].

The main purposes of this paper are:

- to give the construction and existence of all finite limits in $\mathbf{X_2Mod}/\mathbf{R}$.
- to compose the product of the $\mathbf{X_2Mod}/\mathbf{R}$.
- to express the completeness of the $\mathbf{X_2Mod}/\mathbf{R}$.
- to introduce the coequilaser of the $\mathbf{X_2Mod}/\mathbf{R}$.
- to prove the cocompleteness of the category.

We give an example about these structures at the end of this paper.

2. Preliminaries

Definition 2.1 *A crossed module [20], $\partial : R \longrightarrow G$ satisfies two conditions with the action*

$$\begin{aligned} G \times R &\longrightarrow R \\ (g, r) &\mapsto {}^g r \end{aligned}$$

for $g \in G$ and $r \in R$. These conditions are:

- $\varkappa_1) \partial({}^g r) = g\partial(r)g^{-1}$
- $\varkappa_2) {}^{\partial(r)}(r') = rr'r^{-1}$

for $r, r' \in R$ and $g \in G$. With condition $\varkappa_1$ this structure is called a precrossed module. The condition $\varkappa_2$ is known as the peiffer condition. We will denote this category with the crossed module is denoted by **XMod** *and a crossed module $\partial : R \longrightarrow G$ with triple (R, G, ∂).*

Examples:

i) Let $i : R \hookrightarrow G$ be the inclusion map and R be a normal subgroup of G, (R, G, i) is a crossed module. Also, if $\delta : S \longrightarrow G$ crossed module, then δS is a normal subgroup in G.

ii) For any group R, $\partial : R \longrightarrow Aut(R)$ is a crossed module. $Aut(R)$ corresponds to the inner automorphisms.

Definition 2.2 *A 2-crossed module [11] of groups is complex*

$$L \xrightarrow{\partial_2} M \xrightarrow{\partial_1} N$$

of groups with the action of N on L and M. If the mapping

$$\{-,-\} : M \times M \longrightarrow L$$

known as the peiffer lifting satisfies the following conditions,

- $\partial_2\{m_1, m_2\} = {m_2}^{\partial_1(m_1)}(m_1 m_2^{-1}(m_1)^{-1})$
- $\{\partial_2(l_1), \partial_2(l_2)\} = [l_2, l_1]$
- $\{m_1 m_2, m_3\} = \{m_2, m_3\}^{\partial_1(m_1)}\{m_1, m_2 m_3(m_2)^{-1}\}$

  $\{m_1, m_2 m_3\} = \{m_1, m_2\}^{m_1 m_2 (m_1)^{-1}}\{m_1, m_3\}$
- $\{m_1, \partial_2(l_1)\}\{\partial_2(l_1), m_1\} = {l_1}^{\partial_1(m_1)}{l_1}^{-1}$
- ${}^n\{m_1, m_2\} = \{{}^n m_1, {}^n m_2\}$

  for $m_1, m_2, m_3 \in M, l_1, l_2 \in L, n \in N$.

Let $(L_1, M_1, N_1, \partial_2, \partial_1, \{-,-\}_1)$ *and* $(L_2, M_2, N_2, \delta_2, \delta_1, \{-,-\}_2)$ *be 2-crossed modules.* $f = (f_1, f_2, f_3) : (L_1, M_1, N_1) \longrightarrow (L_2, M_2, N_2)$ *is a 2-crossed module morphism if the diagram*

$$\begin{array}{ccccccc} M_1 \times M_1 & \xrightarrow{\{-,-\}_1} & L_1 & \xrightarrow{\partial_2} & M_1 & \xrightarrow{\partial_1} & N_1 \\ \downarrow{\scriptstyle f_2 \times f_2} & & \downarrow{\scriptstyle f_1} & & \downarrow{\scriptstyle f_2} & & \downarrow{\scriptstyle f_3} \\ M_2 \times M_2 & \xrightarrow[\{-,-\}_2]{} & L_2 & \xrightarrow[\delta_2]{} & M_2 & \xrightarrow[\delta_1]{} & N_2 \end{array}$$

is commutative and

$$\begin{aligned} f_2({}^{n_1}m_1) &= {}^{f_3(n_1)}f_2(m_2) \\ f_1({}^{n_1}l_1) &= {}^{f_3(n_1)}f_1(l_1) \end{aligned}$$

for $n_1 \in N_1, m_1 \in M_1$ *and* $l_1 \in L_1$.

3. Finite limits in $\mathbf{X_2Mod/R}$

We give the construction and the existence of all finite limits in $\mathbf{X_2Mod/R}$. In particular for two 2-crossed R-modules we construct their product to use this to obtain finite limits in $\mathbf{X_2Mod/R}$.

Proposition 3.1 *Equaliser object exists in* $\mathbf{X_2Mod/R}$.

Proof Let $(\alpha, \beta) : (A_2, A_1, R, \partial_2, \partial_1) \to (B_2, B_1, R, \delta_2, \delta_1)$ be morphisms in $\mathbf{X_2Mod/R}$ as given with the diagram

$$\begin{array}{ccccc} A_2 & \xrightarrow{\partial_2} & A_1 & \xrightarrow{\partial_1} & R \\ {\scriptstyle\beta_2}\downarrow\downarrow{\scriptstyle\alpha_2} & & {\scriptstyle\beta_1}\downarrow\downarrow{\scriptstyle\alpha_1} & & \| \\ B_2 & \xrightarrow[\delta_2]{} & B_1 & \xrightarrow[\delta_1]{} & R \end{array}$$

where $\alpha = (\alpha_1, \alpha_2)$ and $\beta = (\beta_1, \beta_2)$. Define

$$C_i = \{a_i \in A_i : \alpha(a_i) = \beta(a_i)\}$$


for $i = 1, 2$. With induced morphisms $\varepsilon_1, \varepsilon_2$ via ∂_1 and ∂_2, $(C_2, C_1, R, \varepsilon_2, \varepsilon_1)$ becomes a sub 2-crossed R-module of $(A_2, A_1, R, \partial_2, \partial_1)$. The inclusion $i = (i_2, i_1) : (C_2, C_1, R, \varepsilon_2, \varepsilon_1) \to (A_2, A_1, R, \partial_2, \partial_1)$ is a 2-crossed R-module morphism and satisfies

$$(\alpha_k i_k)(c_k) = \alpha_k(i_k(c_k)) = \beta_k(i_k(c_k)) = (\beta_k i_k)(c_k) \quad , k = 1, 2$$

for all $c_1 \in C_1$ and $c_2 \in C_2$. If $i' = (i'_2, i'_1) : (C'_2, C'_1, R, \varepsilon'_2, \varepsilon'_1) \to (A_2, A_1, R, \partial_2, \partial_1)$ is another 2-crossed R-module morphism such that

$$(\alpha_k i'_k)(c'_k) = (\beta_k i'_k)(c'_k) \quad , k = 1, 2$$

for all $c'_1 \in C'_1$ and $c'_2 \in C'_2$. Then from the definition of C_1 and C_2 we get $i'_2(c'_2) \in C_2$ and $i'_1(c'_1) \in C_1$. Thus we can define $\theta = (\theta_2, \theta_1) : (C'_2, C'_1, \varepsilon') \to (C_2, C_1, \varepsilon)$ as $\theta_k(c'_k) = i'_k(c_k)$ for $k = 1, 2$. Since (i'_2, i'_1) is a 2-crossed R-module morphism, (θ_2, θ_1) is the unique morphism making the diagram



commutative. Thus the morphism (i_2, i_1) is the equalizer of (α, β). □

Proposition 3.2 *Pullback object exists in* $\mathbf{X_2Mod/R}$.

Proof Let $(\alpha_2, \alpha_1) : (A_2, A_1, R, \partial_2, \partial_1) \to (B_2, B_1, R, \delta_2, \delta_1)$ and $(\beta_2, \beta_1) : (C_2, C_1, R, \varepsilon_2, \varepsilon_1) \to (B_2, B_1, R, \delta_2, \delta_1)$ be two morphisms of 2-crossed R-modules where

$$\varkappa_1 : A_2 \xrightarrow{\partial_2} A_1 \xrightarrow{\partial_1} R$$

$$\varkappa_2 : B_2 \xrightarrow{\delta_2} B_1 \xrightarrow{\delta_1} R$$

$$\varkappa_3 : C_2 \xrightarrow{\varepsilon_2} C_1 \xrightarrow{\varepsilon_1} R$$

are 2-crossed R-modules with peiffer liftings $\{_,_\}_C : C_1 \times C_1 \to C_2$, $\{_,_\}_B : B_1 \times B_1 \to B_2$ and $\{_,_\}_A : A_1 \times A_1 \to A_2$. Define:


$$P_k = \{(a_k, c_k) : \alpha_k(a_k) = \beta_k(c_k)\} \subset A_k \times C_k, \qquad k = 1, 2$$

$$p_1 : P_1 \longrightarrow B_1, \quad (a_1, c_1) \mapsto \alpha_1(a_1) = \beta_1(c_1)$$

and

$$\omega : P_2 \longrightarrow P_1, \quad (a_2, c_2) \mapsto (\partial_2(a_2), \varepsilon_2(c_2)).$$

Then we have the following diagram.



Since

$$\begin{aligned}
\delta_1 p_1({}^r(a_1, c_1)) &= \delta_1 p_1(({}^r a_1, {}^r c_1)) \\
&= \delta_1(\alpha_1({}^r a_1)) \\
&= \delta_1(\alpha_1({}^r c_1)) \\
&= r\delta_1(\alpha_1(c_1))r^{-1} \\
&= r(\delta_1 p_1(a_1, c_1))r^{-1}
\end{aligned}$$

for all $r \in R$ and $(a_1, c_1) \in P_1$, the map $\delta_1 p_1$ is a precrossed module. Next, we will show that

$$\varkappa' : P_2 \xrightarrow{\omega} P_1 \xrightarrow{\delta_1 p_1} R$$

is an object in $\mathbf{X_2Mod/R}$ where peiffer lifting $\{_,_\}_P : P_1 \times P_1 \to P_2$ is defined by $[(a_1, c_1),\ (a_1', c_1')] \mapsto (\{a_1, a_1'\}_A, \{c_1, c_1'\}_C)$ for all $(a_1, c_1),\ (a_1', c_1') \in P_1$.

1) ω and $\delta_1 p_1$ are R-equivariant. R acts on itself by conjugation.

2) For all $(a_1, c_1),\ (a_1', c_1') \in P_1$,

$$\begin{aligned}
\omega\{(a_1, c_1),\ (a_1', c_1')\}_P &= \omega(\{a_1, a_1'\}_A, \{c_1, c_1'\}_C) \\
&= [\partial_2\{a_1, a_1'\}_A, \varepsilon_2\{c_1, c_1'\}_C] \\
&= \left({}^{\partial_1(a_1)} a_1' a_1 (a_1')^{-1} (a_1)^{-1}, {}^{\varepsilon_1(c_1)} c_1' c_1 (c_1')^{-1} (c_1)^{-1}\right) \\
&= \left({}^{\delta_1\alpha_1(a_1, c_1)} a_1' a_1 (a_1')^{-1} (a_1)^{-1}, {}^{\delta_1\beta_1(a_1, c_1)} c_1' c_1 (c_1')^{-1} (c_1)^{-1}\right) \\
&= \left({}^{\delta_1\alpha_1(a_1, c_1)} (a_1', c_1') (a_1, c_1) (a_1', c_1')^{-1} (a_1, c_1)^{-1}\right)
\end{aligned}$$

3) For all $(a_2, c_2),\ (a_2', c_2') \in P_2$.

$$\begin{aligned}
\{\omega(a_2, c_2), \omega(a_2', c_2')\}_P &= \{(\partial_2(a_2), \varepsilon_2(c_2)), (\partial_2(a_2'), \varepsilon_2(c_2'))\}_P \\
&= (\{\partial_2(a_2), \varepsilon_2(c_2)\}_A, \{\partial_2(a_2'), \varepsilon_2(c_2')\}_C) \\
&= [a_2, a_2'][c_2, c_2'] \\
&= [(a_2, c_2), (a_2', c_2')]
\end{aligned}$$

4) For all $(a_2,c_2)\in P_2,\ (a_1,c_1)\in P_1$.

$$\begin{aligned}\{(a_1,c_1),\omega(a_2,c_2)\}_P\{\omega(a_2,c_2),(a_1,c_1)\}_P &= \{(a_1,c_1),(\partial_2(a_2),\varepsilon_2(c_2))\}_P\{((\partial_2(a_2),\varepsilon_2(c_2)),(a_1,c_1))\}_P\\ &=(\{a_1,\partial_2(a_2)\}_A,\{c_1,\varepsilon_2(c_2)\}_C)(\{\partial_2(a_2),a_1\}_A,\{\varepsilon_2(c_2),c_1\}_C)\\ &=(\{a_1,\partial_2(a_2)\}_A\{\partial_2(a_2),a_1\}_A,\{c_1,\varepsilon_2(c_2)\}_C\{\varepsilon_2(c_2),c_1\}_C)\\ &={}^{(\partial_1(a_1)}a_2\,(a_2)^{-1},{}^{\varepsilon_1(c_1)}c_2\,(c_2)^{-1})\\ &={}^{(\delta_1\alpha_1(a_1,c_1)}a_2\,(a_2)^{-1},{}^{\delta_1\beta_1(a_1,c_1)}c_2\,(c_2)^{-1})\\ &={}^{\delta_1\alpha_1(a_1,c_1)}\,(a_2,c_2)(a_2,c_2)^{-1}\end{aligned}$$

5) For $(a_0,c_0),(a_1,c_1),(a_2,c_2)\in P_1$

i)

$$\{(a_0,c_0),(a_1,c_1)(a_2,c_2)\}_P=\{(a_0,c_0),(a_1,c_1)\}_P^{(a_0,c_0)(a_1,c_1)(a_0,c_0)^{-1}}\{(a_0,c_0),(a_2,c_2)\}_P$$

ii)


$$\{(a_0,c_0)(a_1,c_1),(a_2,c_2)\}_P={}^{\delta_1 p_1}\{((a_1,c_1),(a_2,c_2)\}_P\left\{(a_0,c_0),(a_1,c_1)(a_2,c_2)(a_1,c_1)^{-1}\right\}_P$$

proof left to the reader as an exercise.

6) For $r\in R$ and $(a_0,c_0),(a_1,c_1)\in P_1$

$$\begin{aligned}{}^r\{(a_0,c_0),(a_1,c_1)\}_P &={}^r(\{a_0,a_1\}_A,\{c_0,c_1\}_C)\\ &=({}^r\{a_0,a_1\}_A,{}^r\{c_0,c_1\}_C)\\ &=(\{{}^ra_0,{}^ra_1\}_A,\{{}^rc_0,{}^rc_1\}_C)\\ &=\{({}^ra_0,{}^rc_0),({}^ra_1,{}^rc_1)\}_P\\ &=\{{}^r(a_0,c_0),{}^r(a_1,c_1)\}_P\end{aligned}$$

With the induced morphisms $\pi_{1,2}:\varkappa'\to\varkappa_1$ and projection morphisms $q_{1,2}:\varkappa'\to\varkappa_3$ we have the following commutative diagram




Here $\alpha\pi=\beta q$, the 2-crossed R-module morphisms commute and the morphisms π and q satisfy the universal property. Let $(\pi_1',\pi_2'):\varkappa'\to\varkappa_1$ and $(q_1',q_2'):\varkappa'\to\varkappa_3$ be any morphisms in $\mathbf{X_2Mod/R}$ such that $\alpha\pi'=\beta q'$

and

$$\varkappa'' : P_2' \to P_1' \to R$$

then there is a unique morphism $(h_1, h_2) : \varkappa'' \to \varkappa'$ given by $h_k(p_k') = (\pi_k'(p_k'), q_k'(p_k'))$ for $k = 1, 2$ and $p_1' \in P_1, p_2' \in P_2$ such that the diagram



commutes. This shows that pullback object exists in $\mathbf{X_2Mod/R}$. □

Proposition 3.3 $\mathbf{X_2Mod/R}$ *has finite products.*

Proof For two 2-crossed R-modules say $\varkappa_1$ and $\varkappa_2$, the product $\varkappa_1 \sqcap \varkappa_2$ will be the pullback over the terminal object $\varkappa_t$ where


$$\varkappa_1 : A_2 \xrightarrow{\partial_2} A_1 \xrightarrow{\partial_1} R$$

$$\varkappa_2 : B_2 \xrightarrow{\delta_2} B_1 \xrightarrow{\delta_1} R$$

$$\varkappa_t : C \xrightarrow{\sigma_2} R \xrightarrow{Id} R$$

with peiffer liftings $\{_,_\}_C : R \times R \to C$, $\{_,_\}_B : B_1 \times B_1 \to B_2$ and $\{_,_\}_A : A_1 \times A_1 \to A_2$ and projection maps $\pi_1 : A_2 \sqcap B_2 \longrightarrow A_2, \pi_2 : A_2 \sqcap B_2 \longrightarrow B_2, \pi_1' : A_1 \sqcap B_1 \longrightarrow A_1$ and $\pi_2' : A_1 \sqcap B_1 \longrightarrow B_1$ the

diagram



is commutative. For all $a_k \in A_k, b_k \in B_k$, $k = 1, 2$; $\vartheta_1 : A_1 \sqcap B_1 \to R$ and $\vartheta_2 : A_2 \sqcap B_2 \to A_1 \sqcap B_1$ is given with

$$\vartheta_1 (a_1, b_1) = \partial_1 \pi_1 (a_1, b_1) = \delta_1 \pi_2 (a_1, b_1)$$

$$\vartheta_2 (a_2, b_2) = (\partial_2 (a_2), \delta_2 (b_2))$$

with peiffer lifting $\{_, _\}_\sqcap : (A_1 \sqcap B_1) \times (A_1 \sqcap B_1) \to A_2 \sqcap B_2$

$$\{(a_1, b_1), (a_1', b_1')\}_\sqcap = (\{a_1, a_1'\}_A, \{b_1, b_1'\}_B)$$

we get a 2-crossed R-module $A_2 \sqcap B_2 \to A_1 \sqcap B_1 \to R$. Using induction $\mathbf{X_2Mod/R}$ has finite products.


Conclusion 3.4 *Since* $\mathbf{X_2Mod/R}$ *has finite products and equalisers,* $\mathbf{X_2Mod/R}$ *has a limit for any functor from a finite category to* $\mathbf{X_2Mod/R}$*. Having all finite limits we say that* $\mathbf{X_2Mod/R}$ *is finitely complete.*

□


4. Finite colimits in $\mathbf{X_2Mod/R}$

Proposition 4.1 *In* $\mathbf{X_2Mod/R}$ *every pair of morphisms with common domain and codomain has a coequaliser.*

Proof Let $(\alpha, \beta) : \varkappa_1 \to \varkappa_2$ be two 2-crossed R-module morphisms as given



Let N_k be a normal subgroup of B_k generated by elements of the form $\alpha_k(a_k) - \beta_k(a_k)$, for $k = 1, 2$ and for all $a_k \in A_k$. Taking $\bar{B}_2 = B_2/N_2$, $\bar{B}_1 = B_1/N_1$ and defining $\bar{\delta}_1 : \bar{B}_1 \to R$, $(b_1 + N_1) \mapsto \delta_1 (b_1)$; $\bar{\delta}_2 : \bar{B}_2 \to \bar{B}_1$, $(b_2 + N_2) \mapsto \delta_2 (b_2) + N_2$ with peiffer lifting $\{_, _\}_N : \bar{B}_1 \times \bar{B}_1 \to \bar{B}_2$, $[(b_1 + N_1), (b_1' + N_1)] \mapsto \{b_1, b_1'\}_B + N_2$, $\bar{\varkappa} = \left(\bar{B}_2, \bar{B}_1, R, \bar{\delta}_2, \bar{\delta}_1\right)$ becomes a 2-crossed R-module and the induced morphism $p{:}\varkappa_2 \to \bar{\varkappa}$ is a 2-crossed R-module morphism such that the diagram




is commutative. Suppose there exists another 2-crossed module morphism $p' : \varkappa_2 \to \varkappa'$ then there is a unique morphism in $\mathbf{X_2Mod/R}$ say $\varphi : \bar{\varkappa} \to \varkappa'$ in which

$$\varphi_k(b_k + N_k) = p'_k(b_k)$$

satisfying $\varphi_k p_k = p'_k$ for $k = 1, 2$ (where $\overline{\varkappa} : \overline{B_2} \xrightarrow{\overline{\delta_2}} \overline{B_1} \xrightarrow{\overline{\delta_1}} R$). Then the morphism $p = (p_1, p_2)$ is universal making the following diagram commutative.

Thus p is the coequaliser of α and β



Let

$$\varkappa_1 : A_2 \xrightarrow{\partial_2} A_1 \xrightarrow{\partial_1} R$$

$$\varkappa_2 : B_2 \xrightarrow{\delta_2} B_1 \xrightarrow{\delta_1} R$$

be two 2-crossed R-modules. Since B_1 acts on A_1 via δ_1 we can form the semidirect product $B_1 \ltimes A_1$ with

$$(b_1, a_1)(b'_1, a'_1) = \left(b_1 \left(b'_1\right)^{\delta_1\left(b'_1\right)}, a'_1\right)$$

and

$$^r(b_1, a_1) = (^r b_1, ^r a_1)$$

for $(b_1,a_1),(b_1',a_1')\in B_1\ltimes A_1$ and $r\in R$. We get injections $i_1:B_1\to B_1\ltimes A_1$, $j_1:A_1\to B_1\ltimes A_1$ and define $\sigma_1:B_1\ltimes A_1\to R$ as $\sigma_1(b_1,a_1)=\delta_1(b_1)+\partial_1(a_1)$. Let N_1 be the normal subgroup of $B_1\ltimes A_1$ generated by the elements

$$(b_1,a_1)(b_2,a_2)-\sigma_1(b_1,a_1)\cdot(b_2,a_2)$$

$$(b_1,a_1)(b_2,a_2)-(b_1,a_1)\cdot\sigma_1(b_2,a_2)$$

Thus the factor group $B_1\ltimes A_1/N_1$ can be defined with the induced morphism $B_1\ltimes A_1/N_1\to R$ as $\bar{\sigma}_1[(b_1,a_1)+N_1]=\sigma_1(b_1,a_1)$. Clearly $\bar{\sigma}_1$ is a precrossed module. Furthermore B_2 acts on A_2 via $\delta_2\delta_1$ and $\delta^2=0$, the semidirect product of A_2 and B_2 is the direct product. With injections $i_2:B_2\to B_2\ltimes A_2$, $j_1:A_2\to B_2\ltimes A_2$ we define $\sigma_2:B_2\ltimes A_2\to B_1\ltimes A_1$ as $\sigma_2(b_2,a_2)=[\delta_2(b_2),\partial_2(a_2)]$. With the same method we construct $B_1\ltimes A_1/N_1$, we can form $B_2\ltimes A_2/N_2$ and the morphism $\bar{\sigma}_2[(b_2,a_2)+N_2]=[\delta_2(b_2),\partial_2(a_2)]+N_1$ and the action of $B_1\ltimes A_1/N_1$ on $B_2\ltimes A_2/N_2$ is given via σ_1. □

Proposition 4.2 *$(B_2\ltimes A_2)/N_2\to(B_1\ltimes A_1)/N_1\to R$ is a 2-crossed R-module.*

Proof First we define peiffer lifting $\{_,_\}_{\bar{N}}:(B_1\ltimes A_1/N_1)\times(B_1\ltimes A_1/N_1)\to B_2\ltimes A_2/N_2$ as $[(b_1,a_1)+N_1,(b_1',a_1')+N_1]\mapsto(\{b_1,b_1'\}_B+N_2,\{a_1,a_1'\}_A+N_2)$ for $(b_1,a_1)+N_1,(b_1',a_1')+N_1\in(B_1\ltimes A_1/N_1)$.

1) $\bar{\sigma}_2$ and $\bar{\sigma}_1$ are R-equivariant. R acts on itself by conjugation.

2) For $(b_1,a_1)+N_1,(b_1',a_1')+N_1\in(B_1\ltimes A_1/N_1)$

$$\begin{aligned}
\bar{\sigma}_2\{(b_1,a_1)+N_1,(b_1',a_1')+N_1\}_N &= \bar{\sigma}_2(\{b_1,b_1'\}_B+N_2,\{a_1,a_1'\}_A+N_2)\\
&= \left[\bar{\delta}_2\{b_1,b_1'\}_B,\bar{\partial}_2\{a_1,a_1'\}_A\right]\\
&= ({}^{\bar{\delta}_1(b_1+N_1)}(b_1'+N_1)(b_1+N_1)(b_1'+N_1)^{-1}(b_1+N_1)^{-1},\\
&\quad {}^{\bar{\partial}_1(a_1+N_1)}(a_1'+N_1)(a_1+N_1)(a_1'+N_1)^{-1}(a_1+N_1)^{-1})\\
&= ({}^{\bar{\sigma}_1[(b_1,a_1)+N_1]}(b_1'+N_1)(b_1+N_1)(b_1'+N_1)^{-1}(b_1+N_1)^{-1},\\
&\quad {}^{\bar{\sigma}_1[(b_1,a_1)+N_1]}(a_1'+N_1)(a_1+N_1)(a_1'+N_1)^{-1}(a_1+N_1)^{-1})\\
&= {}^{\bar{\sigma}_1[(b_1,a_1)+N_1]}((b_1',a_1')+N_1)((b_1,a_1)+N_1)\\
&\quad ((b_1',a_1')+N_1)^{-1}((b_1,a_1)+N_1)^{-1}
\end{aligned}$$

3) For $(b_2,a_2)+N_2,(b_2',a_2')+N_2\in(B_2\ltimes A_2/N_2)$

$$\begin{aligned}
\{\bar{\sigma}_2(b_2,a_2)+N_2,\bar{\sigma}_2(b_2',a_2')+N_2\}_N &= \{(\delta_2(b_2),\partial_2(a_2))+N_1,(\delta_2(b_2'),\partial_2(a_2'))+N_1\}_P\\
&= (\{\delta_2(b_2),\delta_2(b_2')\}_B+N_2,\{\partial_2(a_2),\partial_2(a_2')\}_A+N_2)\\
&= ([b_2,b_2']+N_2,[a_2,a_2']+N_2)\\
&= [(b_2,a_2),(b_2',a_2')]+N_2
\end{aligned}$$

4) For $(b_2,a_2)+N_2\in(B_2\ltimes A_2/N_2),(b_1,a_1)+N_1\in(B_1\ltimes A_1/N_1)$,

$$\begin{aligned}
&\{(b_1,a_1)+N_1, \bar{\sigma}_2(b_2,a_2)+N_2\}_N \{\bar{\sigma}_2(b_2,a_2)+N_2, (b_1,a_1)+N_1\}_N \\
&= (\{b_1+N_1, \delta_2 b_2+N_2\}_B \{\delta_2 b_2+N_2, b_1+N_1\}_B, \{a_1+N_1, \partial_2 a_2+N_2\}_A \{\partial_2 a_2+N_2, a_1+N_1\}_A) \\
&= ((b_2+N_2)^{\delta_1 b_1+N_1}(b_2+N_2)^{-1}, (a_2+N_2)^{\partial_1 a_1+N_1}(a_2+N_2)^{-1}) \\
&= ((b_2+N_2)^{\bar{\delta}_1(b_1,a_1)+N_1}(b_2+N_2)^{-1}, (a_2+N_2)^{\bar{\partial}_1(b_1,a_1)+N_1}(a_2+N_2)^{-1}) \\
&= ((b_2,a_2)+N_2)^{\bar{\sigma}_1(b_1,a_1)+N_1}((b_2,a_2)+N_2)^{-1}
\end{aligned}$$

5) For $(b_k,a_k)+N_1 \in (B_1 \ltimes A_1/N_1),\ k=0,1,2$

$$\begin{aligned}
i.\ \{(b_0,a_0)+N_1,(b_1,a_1)(b_2,a_2)+N_1\}_N &= \{(b_0,a_0)+N_1,(b_1,a_1)+N_1\}_N^{(b_0,a_0)(b_1,a_1)(b_0,a_0)^{-1}} \\
&\quad \{(b_0,a_0)+N_1,(b_2,a_2)+N_1\}_N \\
ii.\ \{(b_0,a_0)(b_1,a_1)+N_1,(b_2,a_2)+N_1\}_N &= {}^{\bar{\sigma}(b_0,a_0)+N_1}\{(b_1,a_1)+N_1,(b_2,a_2)+N_1\}_N \\
&\quad \left\{(b_0,a_0)+N_1,(b_1,a_1)(b_2,a_2)(b_1,a_1)^{-1}\right\}_N \quad \text{left to the reader.}
\end{aligned}$$

6) For $r \in R$ and $(b_0,a_0)+N_1,(b_1,a_1)+N_1 \in (B_1 \ltimes A_1/N_1)$

$$\begin{aligned}
{}^r\{(b_0,a_0)+N_1,(b_1,a_1)+N_1\}_N &= {}^r(\{b_0,b_1\}_B+N_2, \{a_0,a_1\}_A+N_2) \\
&= ({}^r\{b_0,b_1\}_B+N_2, {}^r\{a_0,a_1\}_A+N_2) \\
&= \{{}^r(b_0,a_0)+N_1, {}^r(b_1,a_1)+N_1\}
\end{aligned}$$

□

Theorem 4.3 *The constructed 2-crossed module*

$$\varkappa : (B_2 \times A_2)/N_2 \xrightarrow{\bar{\delta}_2} (B_1 \times A_1)/N_1 \xrightarrow{\bar{\delta}_1} R$$


with the morphisms (i_1,j_1), (i_2,j_2) *is the coproduct of the 2-crossed modules.*

Proof Let $\varkappa_C = (C_2,C_1,R,\partial_2',\partial_1')$ be any 2-crossed R-module and $\alpha : (B_2,B_1,R,\delta_2,\delta_1) \to (C_2,C_1,R,\partial_2',\partial_1')$ and $\beta : (A_2,A_1,R,\partial_2,\partial_1) \to (C_2,C_1,R,\partial_2',\partial_1')$ be 2-crossed R-module morphisms as given by the following diagram.

$$\begin{array}{ccccc}
A_2 & \xrightarrow{\partial_2} & A_1 & \xrightarrow{\partial_1} & R \\
\downarrow{\scriptstyle \beta_2} & & \downarrow{\scriptstyle \beta_1} & & \| \\
C_2 & \xrightarrow{\partial_2'} & C_1 & \xrightarrow{\partial_1'} & R \\
\uparrow{\scriptstyle \alpha_2} & & \uparrow{\scriptstyle \alpha_1} & & \| \\
B_2 & \xrightarrow{\delta_2} & B_1 & \xrightarrow{\delta_1} & R
\end{array}$$

Then there exists a map $h : \varkappa \to \varkappa_C$ given by $h_k[(b_k,a_k)+N_k] = \alpha_k(b_k)\beta_k(a_k)$. $h=(h_1,h_2)$ is a unique

2-crossed R-module morphism making the diagram



commutative. □

The construction of coproducts in $\mathbf{X_2Mod/R}$ induces the functor

$$\circ : \mathbf{X_2Mod/R} \times \mathbf{X_2Mod/R} \to \mathbf{X_2Mod/R}$$

which is left adjoint to the diagonal functor

$$\triangle : \mathbf{X_2Mod/R} \to \mathbf{X_2Mod/R} \times \mathbf{X_2Mod/R}.$$

Conclusion 4.4 $\mathbf{X_2Mod/R}$ *is cocomplete.*

Example 4.5 *Let*

$$\varkappa_1 : B_2 \xrightarrow{\delta_2} B_1 \xrightarrow{\delta_1} R$$

$$\varkappa_2 : A_2 \xrightarrow{\partial_2} A_1 \xrightarrow{\partial_1} R$$

be two 2-crossed R-modules with $\partial_1(A_1) \subset \delta_1(B_1)$ and peiffer liftings $\{_,_\}_A, \{_,_\}_B$. Let $\sigma_1 : \delta_1(B_1) \to B_1$ be an R-equivariant section of δ_1. Then the morphisms

$$\begin{array}{lcll} i_1 & : & B_1 \to B_1 \times A_1/[A_1, B_1]; & b_1 \mapsto (b_1, 0) \\ i_2 & : & B_2 \to B_2 \times B_1; & b_2 \mapsto (b_2, 0) \\ j_1 & : & A_1 \to B_1 \times A_1/[A_1, B_1]; & a_1 \mapsto (\sigma_1(a_1), [a_1]) \\ j_2 & : & A_2 \to B_2 \times A_2; & a_2 \mapsto (0, a_2) \end{array}$$

give a coproduct of 2-crossed modules,

$$\varkappa_1 : B_2 \xrightarrow{\delta_2} B_1 \xrightarrow{\delta_1} R$$

with

$$\begin{aligned}\varepsilon_1 &: B_1 \times (A_1/[A_1,B_1]) \to R; \qquad (b_1,[x]) \mapsto \delta_1(b_1)\\ \varepsilon_2 &: B_2 \times A_2 \to B_1 \times (A_1/[A_1,B_1]); \qquad (b_2,b_1) \mapsto (\delta_2(b_2),0)\end{aligned}$$

and peiffer lifting

$$\begin{aligned}\{_,_\}_{cop} : (B_1 \times (A_1/[A_1,B_1])) \times (B_1 \times (A_1/[A_1,B_1])) &\to B_2 \times A_2\\ [(b_1,[x]),(b_1',[x])] &\mapsto (\{b_1,b_1'\}_B,[\{x_1,x_1'\}_A])\end{aligned}$$

Here $[A_1,B_1]$ *is the normal subgroup of* A_1 *generated by the elements* $a^{-1}a^b$ *for all* $a \in A_1, b \in B_1$.

References

[1] Arslan UE, Akça II, Irmak GO, Avcıoglu O. Fibrations of 2-crossed modules. Mathematical Methods in the Applied Sciences 2019; 42 (16): 5293-5304.

[2] Arvasi Z. Crossed squares and 2-crossed modules of commutative algebras. Theory and Applications of Categories 1997; 3 (7): 160-181.

[3] Arvasi Z, Porter T. Higher dimensional Peiffer elements in simplicial commutative algebras. Theory and Applications of Categories 1997; 3 (1): 1-23.

[4] Arvasi Z, Ulualan E. On algebraic models for homotopy 3-types. Journal of Homotopy and Related Structures 2006; 1 (1): 1-27.

[5] Baues HJ. Combinatorial homotopy and 4-dimensional complexes. Berlin, New York: Walter de Gruyter, 1991.

[6] Brown R, Gilbert ND. Algebraic models of 3-types and automorphism structures for crossed modules. Proceedings of the London Mathematical Society(3) 1989; 59 (1): 51-73.

[7] Brown R, Higgins PJ, Sivera R. Nonabelian algebraic topology: filtered spaces, crossed complexes, cubical higher homotopy groupoids. Zurich, Switzerland: European Mathematical Society, 2011.

[8] Brown R, Sivera R. Algebraic colimit calculations in homotopy theory using fibred and cofibred categories. Theory and Applications of Categories 2009; 22 (8): 222-251.

[9] Carrasco P, Porter T. Coproduct of 2-crossed modules applications to a definition of a tensor product for 2-crossed complexes. Collectanea Mathematica 2016; 67 (3): 485-517.

[10] Casas JM, Ladra M. Colimits in the crossed modules category in Lie algebras. Georgian Mathematical Journal 2000; 7 (3): 461-474.

[11] Conduché D. Modules croisés généralisés de longueur 2. Journal of Pure and Applied Algebra 1984; 34 (2-3): 155-178.

[12] Conduché D. Simplicial crossed modules and mapping cones. Georgian Mathematical Journal 2003; 10 (4): 623–636.

[13] Ellis GJ. Higher dimensional crossed module of algebras. Journal of Pure and Applied Algebra 2003; 52 (3): 277-282.

[14] Ellis GJ. Crossed squares and combinatorial homotopy. Mathematische Zeitschrift 1993; 214 (1): 93-110.

[15] Ellis GJ, Mikhailov R. A colimit classifying spaces. Advances in Mathematics 2010; 223 (6): 2097-2113.

[16] Guin-Waléry D, Loday J-L. Obstructioná l'excision en K-théorie algébrique, in: Algebraic K-Theory. In: Friedlander EM, Stein MR (editor) Algebraic K-Theory Evanston 1980. Berlin, Heidelberg, Germany: Springer, 1981, pp. 179-216.

[17] Grandjéan AR, Vale MJ. 2-Modulos cruzados an la cohomologia de André- Quillen. Memorias de la Real Academia de Ciencias 1986; 22: 1-28.

[18] Mutlu A, Porter T. Freeness conditions for 2-crossed modules and complexes. Theory and Applications of Categories 1998; 4 (8): 174–194.

[19] Shammu NM. Algebraic and categorical structures of category of crossed modules of algebras. PhD, University of Wales, Bangor, United Kingdom, 1992.

[20] Whitehead JHC. Combinatorial homotopy II. Bulletin of the American Mathematical Society 1949; 55 (5): 453-496.

[21] Yılmaz K. 2-Crossed modules over same base. Sakarya University Journal of Science 2017; 21 (6): 1210-1220.