HPR Patients' perspectives, functioning and health (descriptive: qualitative or quantitative)

POS1583-HPR THE EFFECT OF VITAMIN D DEFICIENCY ON EXERCISE CAPACITY, RESPIRATORY MUSCLE STRENGTH, AND PERIPHERAL MUSCLE STRENGTH IN PATIENTS WITH CONNECTIVE TISSUE DISEASES

Keywords: Vitamin D, Systemic sclerosis, Systemic lupus erythematosus

F. Sarı[1], D. Saraç[2], S. Bayram[3], S. Baglan Yentur[4], N. G. Tore[3], D. Oskay[3], A. Tufan[5]. *[1]Bingöl University, Department of Physiotherapy and Rehabilitation, Faculty of Health Sciences, Bingöl, Turkey; [2]Izmir Katip Çelebi University, Department of Physiotherapy and Rehabilitation, Faculty of Health Sciences, İzmir, Turkey; [3]Gazi University, Department of Physiotherapy and Rehabilitation, Faculty of Health Sciences, Ankara, Turkey; [4]Firat University, Department of Physiotherapy and Rehabilitation, Faculty of Health Sciences, Elazığ, Turkey; [5]National Human Genome Research Institute, Inflamatuar Disease Section, Rockville Pike, United States of America*

Background: Vitamin D is one of the fat-soluble vitamins and is also a group of sterols that are hormone and hormone precursors. Vitamin D deficiency and/or insufficiency have been found to be associated with many chronic diseases, including cancers, metabolic syndrome, and autoimmune diseases. Vitamin D is involved in modulation of immune responses in autoimmune disorders including multiple sclerosis, asthma, diabetes mellitus, connective tissue disorders such as rheumatoid arthritis (RA) or systemic lupus erythematosus (SLE). In addition; Previous studies conducted in other patient populations indicate that higher vitamin D levels are related to higher exercise capacity, muscle strength, and better lung function.
Objectives: The purpose was to investigate the effects of vitamin D deficiency on exercise capacity, respiratory muscle strength, and peripheral muscle strength in patients with connective tissue disease (CTD).
Methods: This cross-sectional observational study included CDT (SLE and systemic sclerosis (SSc)) patients and healthy participants. Serum 25(OH)D, calcium, and phosphorus levels were measured and Vitamin D levels 10.0-20.0 ng/mL defined insufficiency and levels <10.0 ng/mL defined deficiency. The exercise capacity, respiratory muscle strength, and peripheral muscle strengths were set as the primary outcomes of the study. Secondary outcomes included evaluation of physical activity, dyspnea, pain, emotional status, fatigue, and quality of life.
Results: There were 23 SLE (36.82±10.06), 21 SSc (44.38±14.63), and 24 healthy controls (37.62±11.74). Compared to healthy controls, CTD groups with vitamin D deficiency had significantly lower respiratory muscle strength (p<0.001), peripheral muscle strength (knee extension and shoulder flexion) (p<0.05), exercise capacity (p<0.05), physical activity (p<0.05), and quality of life (p<0.05) while having significantly more dyspnea (p<0.001), pain (p<0.001), fatigue (p<0.001), anxiety, and depression (p<0.05).
Conclusion: The results of our study indicate that vitamin D deficiency is associated with reduced muscle strenghts and related symptoms and complaints in CTD patients, hence, in order to increase the efficacy of rehabilitation and treatment programs, attention should be paid to vitamin D levels and should be corrected appropriately.

REFERENCES:

[1] Sizar O, Khare S, Goyal A, Givler A. Vitamin D Deficiency. StatPearls. Treasure Island (FL): StatPearls Publishing. Copyright © 2022, StatPearls Publishing LLC.; 2022.
[2] Vaidya B, Nakarmi S. Vitamin D in Rheumatic Diseases: Interpretation and Significance. Fads and Facts about Vitamin D [Internet]. London, United Kingdom: Intech Open; 2019.
[3] Hassanalilou T, Khalili L, Ghavamzadeh S, Shokri A, Payahoo L, Bishak YK (2017) Role of vitamin D deficiency in systemic lupus erythematosus incidence and aggravation. Auto Immun Highlights 9(1):1. https://doi.org/10.1007/s13317-017-0101-x.
[4] Dawson-Hughes B (2017) Vitamin D and muscle function. J Steroid Biochem Mol Biol 173:313-6. https://doi.org/10.1016/j.jsbmb.2017.03.018.
[5] Ahmed B, Nasir K, Mehmood A, Abid MA, Zehra NA, Khan AH, et al. (2021) Effect of physical activity and vitamin D compared with vitamin D alone on muscle strength, back flexibility and aerobic activity in patients with chronic kidney disease: A comparative study from Pakistan. Asia Pac J Clin Nutr 30(4):566-72. https://doi.org/10.6133/apjcn.202112_30(4).0002.

Acknowledgements: NIL.
Disclosure of Interests: None Declared.
DOI: 10.1136/annrheumdis-2023-eular.3544

POS1584-HPR IMPACT OF TELEMEDICINE ON DISEASE ACTIVITY ASSESSMENT: A CASE-CROSS OVER NESTED WITHIN A COHORT OF SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS

Keywords: Systemic lupus erythematosus, Outcome measures, COVID

I. Almaghlouth[1], H. Almalag[2], H. Alzuhair[1], A. Bedaiwi[1], L. Hassen[1], S. Alzomia[1], B. Alanazi[1], S. Alahmari[1], K. N. Asfina[1], N. Khalil[1], M. A. Omair[1], M. Bedaiwi[1]. *[1]King Saud University, Rheumatology Unit, Department of Medicine, Riyadh, Saudi Arabia; [2]King Saud University, Clinical pharmacy, Riyadh, Saudi Arabia*

Background: The utilization of telemedicine has been rapidly growing among patients with rheumatic disease especially following COVID-19 pandemic. The ease and convenience seems to dominate the reasons of such growth. Yet the effect of this approach on patients with systemic lupus erythematosus (SLE) is yet to be revealed.
Objectives: In this study, we aimed to examine the effect of telemedicine in SLE patients on the outcome of disease activity.
Methods: This is a case-cross over study nested within the national prospective cohort of SLE, Saudi Arabia. SLE patients who fulfils the SLICC classification criteria within the period between March 2020 to March 2021 and have been assessed at three-time point with three months' time difference between assessments according to standardized protocol were included. Telemedicine was conducted at first point while in person assessment were used for second and third points. Primary outcome was difference in SLEDAI-2K score. Primary analysis was conducted using GEE model and adjusted for potential confounders including demographics, medications and changes in steroid doses. Several sensitivity analyses were conducted to mitigate selection and time varying confounders.
Results: A total of 92 participants were included. Majority of them were females (88%), with a mean (±SD) age of 36 (±13). mean (±SD) disease activity scores at baseline as follows: SLEDI 5 (±5); SRI 3.8 (±3.5); SDI 1 (±1). Mean difference of SLEDI score of -1.641 (95% CI -2.773 — -0.510), p=0.005* between telemedicine and follow up visits. Adjusted value and mean with mean diffrence in Figure 1. Results were consistent in all sensitivity analyses.
Conclusion: We found that the Telemedicine assessment was associated with much higher disease activity score compared to in person in subsequent assessments which may suggest potentially overestimation of disease activity and later assessment accuracy. Cautious adoption is suggested in SLE patients with active disease.

REFERENCES:

[1] Anderson JT, Bouchacourt LM, Sussman KL, Bright LF, Wilcox GB. Telehealth adoption during the COVID-19 pandemic: A social media textual and network analysis. Digit Health. 2022 Mar 31;8:20552076221090041. doi: 10.1177/20552076221090041. PMID: 35392254; PMCID: PMC8979849.
[2] Tang, W., Inzerillo, S., Weiner, J., Khalili, L., Barasch, J., Gartshteyn, Y., Dall'Era, M., Aranow, C., Mackay, M.C., & Askanase, A.D. (2022). The Impact of Telemedicine on Rheumatology Care. Frontiers in Medicine, 9.




Figure 1 Repeated measure analysis of difference in SLEDAI between the virtual visit and the physical visit taking factors affecting SLEDAI on univariate analysis into consideration (abnormal urine test, fibromyalgia, haemolytic anaemia and mycophenolate use).

Acknowledgements: NIL.
Disclosure of Interests: None Declared.
DOI: 10.1136/annrheumdis-2023-eular.3695

POS1585-HPR "SO, MEN WILL BE ABLE TO RECEIVE #METHOTREXATE FOR LUPUS AND RHEUMATOID ARTHRITIS BUT NOT WOMEN" THE OVERTURN OF ROE V. WADE BEYOND ABORTION

Keywords: Qualitative research methods, Pregnancy and reproduction, Quality of life

J. B. Negrón[1], M. Rodríguez-Malavé[1]. *[1]Instituto de Investigación Social y Sanitaria, Departamento de Investigación, Guayanilla, Puerto Rico*

Ann Rheum Dis: first published as 10.1136/annrheumdis-2023-eular.3544 on 30 May 2023. Downloaded from http://ard.bmj.com/ on February 27, 2024 at Eskisehir Osmangazi University. Protected by copyright.